

Isospin Multiplet Structure in Ultra-Heavy Fermion Bound States

**Pankaj Jain,^a Alan J. Sommerer,^b Douglas W. McKay,^a
J.R. Spence,^b J.P. Vary,^b and Bing-Lin Young,^b**

^a Department of Physics and Astronomy, University of Kansas, Lawrence, Kansas 66045–2151

^b Department of Physics and Astronomy and Ames Laboratory, Iowa State University, Ames, Iowa 50011.

Abstract

The coupled Bethe–Salpeter bound state equations for a $Q\bar{Q}$ system, where $Q = (U, D)$ is a degenerate, fourth generation, super-heavy quark doublet, are solved in several ladder approximation models. The exchanges of gluon, Higgs and Goldstone modes in the standard model are calculated in the ultra-heavy quark limit where weak γ , $W^\pm$ and Z^0 contributions are negligible. A natural $I = 0$ and $I = 1$ multiplet pattern is found, with large splittings occurring between the different weak iso-spin states when M_Q , the quark masses, are larger than values in the range $0.4TeV < M_Q < 0.8TeV$, depending on which model is used.

Consideration of ultra-heavy quark lifetime constraints and $U - D$ mass splitting constraints are reviewed to establish the plausibility of lifetime and mass degeneracy requirements assumed for this paper.

1. Introduction

The prospect of SSC and LHC experiments to probe TeV phenomena has stimulated a wide variety of ideas about the physics at these energies. The standard model itself, expanded to a super-heavy fourth generation, could display some striking new effects. There is a variety of “heavy-neutrino” fourth generation models in the literature,¹ and we adopt the attitude that some model of this type with the Higgs mechanism providing the fermion and gauge boson mass generation represents nature at 0.5-1.0 TeV. The effect that we explore in this paper is the strong Yukawa coupling of the Higgs sector to the fourth generation and the bound state spectrum in the heavy quark-anti-quark system ($Q\bar{Q}$) that is generated by this strong coupling.^{2,3} There have been a number of heavy fermion physics studies since the Higgs mechanism and the standard model were proposed.^{4,5} Like the heavy Higgs limit studies, the indications are that perturbative calculations break down and some new, strong coupling features of the standard model and/or totally new physics should emerge at the 0.5–1.0 TeV energy ranges.

In a previous work,³ we described some new, deep-binding features of the 0^- ground state of a single, super-heavy $Q\bar{Q}$ system that were driven by the dominant, attractive Higgs scalar exchange plus weaker, but in some cases still significant, gluon exchange. The general feature was that a very strong binding compared to QCD alone was generally exhibited for $M_Q > 0.4 - 0.5$ TeV in all of the model calculations that we tested.

In this paper, we expand significantly on this theme by doing a coupled channel analysis of a heavy quark doublet, (U, D) , including the Goldstone boson exchanges as well as the Higgs exchanges, and we discuss the s -channel exchange as well as the t -channel exchange contributions to the Bethe-Salpeter equation kernel. Here we include the results of calculations of the 0^+ and 1^- masses in addition to the 0^- ground state case analyzed previously.

Because of the weak iso-doublet structure of the super-heavy (U, D) system the bound states decomposes naturally into weak iso-triplet and weak iso-singlet states when $M_U \simeq M_D$ and $M_Z \simeq M_W$. We find that the interplay between the scalar Higgs exchange and the (opposite sign) pseudoscalar Goldstone boson (longitudinal W and Z) exchange results in a clear iso-triplet/iso-singlet mass splitting in all of the channels in all model calculations we used. This signature is quite distinct for the fourth generation fermion doublet, since it must be approximately degenerate according to present constraints from W/Z physics.

In the following section, we outline the coupled-channel Bethe-Salpeter formalism used to calculate bound state masses. In Sec. 3 we present results of the calculations in the different approximation schemes used and comment on the main features. In Sec. 4 we show that requiring meta-stability of fourth generation quarks puts only weak constraints on mixing with a lighter generation. We also review the evidence that a fourth generation of quarks, if they exist, would have to be approximately degenerate in mass. In Sec. 5 we summarize and conclude. In Appendix A, we describe the subtraction used to regulate the Bethe-Salpeter equations. In Appendix B, we give a short analysis of the running of the Yukawa coupling and its relevance to the calculations presented.

2. Strong Yukawa Binding of Ultra-Heavy Fermion Weak Doublets: Bethe-Salpeter Equation Formalism Including Goldstone Boson Exchange

If there are fourth and higher generations of fermion weak isospin doublets, the fractional mass-splitting within these ultra-heavy doublets⁶ is constrained by the standard, three generation model's remarkable survival after two decades of stringent experimental tests.⁷ A small mass-splitting, or near degeneracy, could lead to a classic isospin-multiplet pattern for the bound state spectrum, and we show in this section that this is indeed the case when the Yukawa couplings of the Higgs and Goldstone-bosons⁸ are incorporated into the bound state equations – ladder approximation, Bethe-Salpeter equations in our study. We assume that the usual mass generation mechanism operates in the fourth generation sector as well, so the ultra heavy fermion doublets will have strong Yukawa couplings to the standard model Higgs and Goldstone modes. These Yukawa couplings, scalar for the Higgs boson and pseudoscalar for Goldstone bosons, become the dominant ones when the quark masses reach 400–500 GeV (though the pseudoscalar exchanges actually become competitive only in the tight binding, highly relativistic regime. They decouple in the weak-binding limit).

We develop below a set-up for the two, coupled-channel Bethe-Salpeter equations of a quark doublet-antidoublet that interact through gluon and weak-sector Higgs and Goldstone boson exchange.

Let us consider a doublet of heavy quarks, (U, D) , interacting in the Feynman gauge by exchange of gluons, g , a Higgs boson, H , and Goldstone bosons $\chi^\pm, \chi^0$. The photon and $W^\pm$ and Z exchanges are always weak in the gauges chosen, have been checked and found to give insignificant contributions to the binding, and are not included in our development. It is convenient to adopt Feynman gauge in the ladder approximation (perturbative vertices

no crossed graphs), which we will use throughout this work. There are four quark–antiquark channels: $\bar{U}U$, $\bar{U}D$, $\bar{D}U$ and $\bar{D}D$. These channels become coupled when the charged Goldstone boson exchanges are included, as illustrated in Fig. 1, where the Bethe–Salpeter equations are shown graphically. As discussed below, these equations admit a subtraction at fixed q that eliminates the divergent, q -independent, annihilation graphs which enter the scalar and pseudoscalar bound state channels.⁹ The corresponding gluon annihilation graph does not require consideration because we study only color singlet bound states.

The coupled-channel, doublet quark–antiquark bound state equations are shown pictorially in Fig. 1 and displayed in Eqs. (2.1) and (2.2) below. The subtracted, decoupled versions of these equations, are presented later in this section, Eqs. (2.3a and 2.3b). Let us display, in shorthand notation, the structure of the ladder approximation, momentum – space Bethe–Salpeter equations for the bound state amplitudes $\chi(q, P)$ in Feynman gauge.

$\bar{U} - U$ Channel:

$$\begin{aligned}
S^{-1}(q_+) \chi_{UU}(q, P) S^{-1}(q_-) = & \int \frac{d^4 k}{(2\pi)^4} \left[-\gamma_\mu \chi_{UU}(k, P) \gamma_\nu G^{\mu\nu}(k - q) \right. \\
& - \frac{g_2^2}{4} \frac{m^2}{M_W^2} \chi_{UU}(k, P) \frac{i}{(k - q)^2 - M_H^2} + \frac{g_2^2}{4} \frac{m^2}{M_W^2} \gamma_5 \chi_{UU}(k, P) \gamma_5 \frac{i}{(k - q)^2 - M_Z^2} \\
& + \frac{g_2^2}{2} \frac{m^2}{M_W^2} \gamma_5 \chi_{DD}(k, P) \gamma_5 \frac{i}{(k - q)^2 - M_W^2} \\
& \left. + \frac{g_2^2}{4} \frac{m^2}{M_W^2} \frac{i}{P^2 - M_Z^2} \gamma_5 \text{Tr} \left((\chi_{UU} - \chi_{DD}) \gamma_5 \right) + \frac{g_2^2}{4} \frac{m^2}{M_W^2} \frac{i}{P^2 - M_H^2} \text{Tr} (\chi_{UU} + \chi_{DD}) \right]
\end{aligned} \tag{2.1}$$

$\bar{D} - D$ Channel: interchange $U \leftrightarrow D$ in (2.1).

$\bar{U} - D$ Channel:

$$\begin{aligned}
S^{-1}(q_+) \chi_{UD}(q, P) S^{-1}(q_-) = & \int \frac{d^4 k}{(2\pi)^4} \left[-\gamma_\mu \chi_{UD}(k, P) \gamma_\nu G^{\mu\nu}(k - q) \right. \\
& - \frac{g_2^2}{4} \frac{m^2}{M_W^2} \chi_{UD}(k, P) \frac{i}{(k - q)^2 - M_H^2} - \frac{g_2^2}{4} \frac{m^2}{M_W^2} \gamma_5 \chi_{UD}(k, P) \gamma_5 \frac{i}{(k - q)^2 - M_Z^2} \\
& \left. + \frac{g_2^2}{2} \frac{m^2}{M_W^2} \frac{i}{P^2 - M_W^2} \gamma_5 \text{Tr} (\chi_{UD}(k, P) \gamma_5) \right] .
\end{aligned} \tag{2.2}$$

$\bar{D} - U$ Channel: interchange $U \leftrightarrow D$ in Eq. (2.2).

In Eqs. (2.1) and (2.2), $\chi_{UU}(q, P)$ etc., represent the Bethe–Salpeter amplitudes for the $\bar{U}U$ etc., systems with relative 4-momentum q and total 4-momentum P . The quark masses, m , are assumed to be degenerate, $q_\pm \equiv q \pm \frac{P}{2}$ and $S^{-1}(q_+)$ are the inverse quark

propagators with momenta $q_{\pm}$. Here $G^{\mu\nu}(k-q)$ represents the gluon propagator, assumed to behave like $(k-q)^{-2}$ as $|k-q| \rightarrow \infty$. The SU(2) coupling constant is denoted by g_2 .

One may decouple the $\bar{U}U$ and $\bar{D}D$ equations by taking the combinations $(\bar{U}U \pm \bar{D}D)/\sqrt{2}$, and the resulting Bethe–Salpeter equations, subtracted as described in Appendix A, for the bound state wave–functions $\chi_{UD}(q, P)$, $\chi_{-}(q, P)$ and $\chi_{+}(q, P)$ read as follows ($\chi_{DU}(q, P)$ obeys the same equations as χ_{UD}):

$$S^{-1}(q_{+})\chi_{\pm}(q, P)S^{-1}(q_{-}) = \int \frac{d^4k}{(2\pi)^4} \left\{ -\gamma_{\mu}\chi_{\pm}(k, P)\gamma_{\nu}G^{\mu\nu}(k-q) \right. \\ \left. -i\frac{m^2g_2^2}{4M_W^2}\frac{\chi_{\pm}(k, P)}{(k-q)^2 - M_H^2} + \frac{i}{4}\frac{m^2g_2^2}{M_W^2}\frac{\gamma_5\chi_{\pm}(k, P)\gamma_5}{(k-q)^2 - M_Z^2} \mp \frac{i}{2}\frac{m^2g_2^2}{M_W^2}\frac{\gamma_5\chi_{\pm}(k, P)\gamma_5}{(k-q)^2 - M_W^2} \right\} \quad (2.3a)$$

and

$$S^{-1}(q_{+})\chi_{UD}(q, P)S^{-1}(q_{-}) = \int \frac{d^4k}{(2\pi)^4} \left\{ -\gamma_{\mu}\chi_{UD}(k, P)\gamma_{\nu}G^{\mu\nu}(k-q) \right. \\ \left. -i\frac{m^2g_2^2}{4M_W^2}\frac{\chi_{UD}(k, P)}{(k-q)^2 - M_H^2} - i\frac{m^2g_2^2}{4M_W^2}\frac{\gamma_5\chi_{UD}\gamma_5}{(k-q)^2 - M_Z^2} \right\} \quad (2.3b)$$

The quark mass $m = m_U = m_D$ is taken to be the same for U and D . Experimental support for this assumption is reviewed in Sec. 4.

The important features of Eq. (2.3) are

- (i) χ_{-} , χ_{UD} and χ_{DU} obey the *same equation* when $M_Z = M_W$, and they form an isospin triplet of states.
- (ii) The Goldstone boson contributions to this χ_{-} , χ_{UD} , χ_{DU} triplet are, like the gluon and Higgs boson contributions *all attractive* in the *ground state*, 0^{-} system, (see Sec. 3 below)
- (iii) the χ_{+} state obeys a different equation whose Goldstone–boson contributions are all *repulsive* in the *ground state*, 0^{-} system.

Eq. (2.3) and the fact that pseudoscalar exchange becomes comparable to scalar for relativistic systems combine to produce a clear splitting between the singlet and triplet of ground state bosons in the strong binding limit. The triplet is below the singlet in mass for the 0^{-} system. This is very reminiscent of a “ $\pi - \eta$ ” situation, except that this appears to be purely dynamical with no spontaneous symmetry breaking, bound state Goldstone phenomenon at work.

The iso–singlet, iso–triplet situation is reversed in the 0^{+} and 1^{-} spin–parity channels, as we describe in the next section.

3. Results of Relativistic Bound State Calculations of Iso-Singlet and Iso-Triplet Masses

We employ two different approaches to solving the ladder approximation, Bethe–Salpeter equations for the $J^P = 0^-$ ground states of the weak iso-spin singlet and weak iso-spin triplet $Q\bar{Q}$ system. We also report results in the same approximations for the $J^P = 1^-$ and 0^+ states. The principle feature that emerges, as anticipated in the preceding section, is the splitting between the iso-singlet and iso-triplet states due to the pseudoscalar exchange (in Feynman gauge): repulsion in the iso-singlet case and attraction in the iso-triplet case in the ground state, 0^- , channel. The iso-triplet pseudoscalar, ground state mass solutions are found to fall to zero for high enough quark masses in all of the calculations. The reverse is true in the 0^+ and 1^- channels, and the isoscalar masses fall to zero at high enough quark masses in these states.¹⁰

We evaluate the bound state energies (masses) using both the covariant kernel,

$$K(q) = g^2 \frac{1}{q^2 - M^2} \quad , \quad (3.1a)$$

which we refer to as the covariant gauge ladder approximation, and the instantaneous approximation to this kernel

$$K(|\vec{q}|^2) = -g^2 \frac{1}{|\vec{q}|^2 + M^2} \quad . \quad (3.1b)$$

We have suppressed any reference to gauge-dependent spin structure in (3.1), and designate generic coupling constants and masses as g and M , respectively. The instantaneous approximation, so called because the coordinate space potentials are instantaneous, reduces the problem to three dimensions, yielding what are referred to as Salpeter’s equations. These are coupled equations for positive and negative frequency amplitudes. One can solve the coupled system or set the negative frequency amplitudes to zero and solve the positive frequency problem by itself.

Covariant Gauge Formalism

In the covariant gauge ladder (CGL) formalism, we decompose the 0^- Bethe–Salpeter amplitudes into four independent spinor functions:

$$\chi_P(P, q) = \gamma_5 \left[\chi_0 + \not{P} \chi_1 + \not{\not{q}} \chi_2 + [\not{q}, \not{P}] \chi_3 \right] \quad . \quad (3.2)$$

In our calculation, only χ_0 and χ_1 need to be retained, and the weak coupling relation $\chi_- = -\chi_0/2m$ can be employed to reduce the system of equations to a single equation for

χ_0 . In Euclidean variables, we have

$$\begin{aligned}\chi_0(P, q) = & \frac{4\alpha_s}{3\pi^3} \frac{(q^2 + m^2)}{D} \int d^4k \frac{\chi_0}{(k - q)^2} \\ & + \frac{m^2 g_2^2}{4M_W^2} \frac{1}{D} \left(q^2 + \frac{3M_B^2}{4} + m^2 \right) \int \frac{d^4k}{(2\pi)^4} \frac{\chi_0}{(k - q)^2 + M_H^2} \quad , \quad (3.3) \\ & + \frac{m^2 g_2^2}{4M_W^2} \frac{F^I}{D} \left(q^2 - \frac{M_B^2}{4} + m^2 \right) \int \frac{d^4k}{(2\pi)^4} \frac{\chi_0}{(k - q)^2 + M_Z^2}\end{aligned}$$

where $M_Z = M_W = 90 \text{ GeV}$ has been adopted since including weak boson mass difference affects answers only in the third decimal place, $D \equiv D(q^2, M_B^2, \cos\theta) = (-q^2 + \frac{M_B^2}{4} - m^2)^2 + M_B^2 q^2 \cos^2 \theta$, $M_B^2 = -P^2$ is the mass-squared of the bound state, θ is the angle between P and q and $F^I = +1, -3$ for iso-spin $I = 1, 0$ respectively. As mentioned in the preceding section, the sign difference in the last term between the $I = 1$ and $I = 0$ cases produces the extra attraction and repulsion compared to the purely attractive gluon and Higgs potentials. The method of solution is discussed in Ref. 3 in some detail. The resulting bound state mass values as a function of quark mass, m , are shown in Fig. 2 in the curves labeled S1 and T1. In this calculation, the iso-singlet (S1) state's binding energy steadily increases, but the bound state mass never sinks to zero in the fermion mass range we have investigated. The more tightly bound triplet state (T1) achieves zero mass when the quark mass reaches about 900 GeV. Above 750 GeV quark mass, the splitting between the different bound state masses becomes quite pronounced. The 0^+ and 1^- calculations proceed in a similar fashion with the isosinglet being more tightly bound than the isotriplet. The details are not presented here, since they do not illuminate the discussion.

Salpeter Equation Formalism

Making the approximation 3.1b, integrating over q_0 in the Bethe-Salpeter equation, and projecting out the 0^- channel amplitude, one obtains coupled equations for the positive and negative frequency amplitudes, $\chi_{\pm}(q)$, where $q \equiv |\vec{q}|$. The equations read

$$\begin{aligned}(E - 2\omega)\chi_+(q) &= \frac{1}{\pi q} \int dq' q' (V_+ \chi_+ + V_- \chi_-) \\ (E + 2\omega)\chi_-(q) &= \frac{-1}{\pi q} \int dq' q' (V_- \chi_+ + V_+ \chi_-)\end{aligned}\quad (3.4)$$

where $\omega = \sqrt{q^2 + m^2}$, E is the bound state eigenvalue and

$$\begin{aligned} V_{\pm} = & C_V Q_0(Z_V) \frac{2\omega\omega' \mp m^2}{\omega\omega'} \\ & + C_{PS} \left\{ Q_0(Z_{PS}) \frac{\omega\omega' \mp m^2}{\omega\omega'} + \left[Z_{PS} Q_0(Z_{PS}) - 1 \right] \frac{qq'}{\omega\omega'} \right\} \\ & + C_S \left\{ Q_0(Z_S) \frac{\omega\omega' \pm m^2}{\omega\omega'} + \left[Z_S Q_0(Z_S) - 1 \right] \frac{qq'}{\omega\omega'} \right\} . \end{aligned}$$

The coefficients C_V and variables Z_V etc. are summarized in Table 1. Details on the method of solution are given in Ref. 3.

The 0^- , ground state masses as a function of quark mass is shown for the positive frequency only case in Fig. 2, curves S2 and T2. Again the splitting between isosinglet (S2) and isotriplet (T2) bound state masses is striking for the heavy quark mass region, and the iso-triplet system becomes ultrarelativistic in the region above $m = 750 GeV$, producing a zero mass bound state at $m \simeq 1100 GeV$.

The solution for the positive plus negative, fully coupled, system is much more tightly bound as is seen in Fig. 2, curves S3 and T3, where a dramatic departure between iso-triplet (T3) and iso-singlet (S3) masses sets in already at $m = 400 GeV$, and the iso-triplet mass plunges to zero at $m = 520 GeV$. Figure 2 also shows the feature of the instantaneous approximation that differs from the CGL approximation, namely the turn-over (and eventual fall to zero) of the iso-singlet bound state mass as a function of the quark mass. This effect does not yet appear in Fig. 2 for the iso-singlet positive-frequency-only case, but it does occur at a quark mass above 1.20 TeV for this case as well.

Focusing on the iso-triplet ground state mass values vs. quark mass curves in Fig. 2 for the three solutions which were discussed above, we see that the general features agree though the different relativistic bound-state approximations produce different bound state mass values for a given quark mass. One can conclude that the Goldstone boson Higgs exchange plays an important role in the calculations, that deep binding at or above 500 GeV quark mass occurs, and that a dramatic weak iso-spin mass splitting is produced in the $Q\bar{Q}$ spectrum of a degenerate, or nearly so, U, D doublet system.

With the role of iso-singlet and iso-triplet reversed, the features just outlined are present also in the $J^P = 1^-$ and 0^+ bound states, and we display results of calculations for this system in Figs. 3 and 4. The principle distinction between 0^- on one hand and 1^-

and 0^+ on the other is the iso-singlet/triplet splitting reversal and that the latter two are typically less tightly bound.

In the next section we take up the question of ultra-heavy quark decay lifetime and intra-doublet mass splitting constraints.

4. Constraints on Super-Heavy U, D Quark Lifetimes and on $U - D$ Splitting

Two related questions arise when one considers the phenomenology of super-heavy fermions. A crucial consideration for bound state physics is the comparison of quark lifetime to the period of bound state motion. The situation for the heavy top quark case has been considered by a number of authors,¹² and we apply the argument of Strassler and Peskin to the super-heavy, fourth generation in this section. Related to the lifetime question is the $U - D$ mass-splitting question. If the splitting is larger than the W -mass, then the higher mass quark can decay to the lower mass quark by direct W -emission, presumably with no Cabbibo, Kobayashi-Maskawa (C-K-M) suppression. The width would then be too broad to permit formation of narrow bound states containing the heavier quark. We review constraints imposed by measured W and Z -boson properties on the $U - D$ mass splitting, which we would like to take as degenerate to a first approximation.

Lifetime for decay to light quarks

The lighter of the U, D doublet partners must decay to lighter generation quarks, q , and we assume that one transition is dominant. Calling the corresponding C-K-M factor V_{Qq} , we have

$$\Gamma_{Q \rightarrow q} \sim (180) MeV \left[\frac{M_Q}{M_W} \right]^3 |V_{Qq}|^2, \quad (4.1)$$

where $M_Q \gg M_q, M_W$ is assumed. To make a conservative estimate, we will include only the gluon binding in estimating the time needed for bound state formation in a non-relativistic, weak binding approximation. Including the strong binding, Yukawa interactions will only improve the prospects for bound state formation. For pure QCD coupling, the characteristic radius is $a_0 = (\frac{4}{3}\alpha_s M_Q/2)^{-1}$ where $\frac{4}{3}\alpha_s$ is the effective QCD coupling strength for the problem (with a characteristic velocity $v_n \sim (4/3)\alpha_s/n$ for the n^{th} radial excitation). Taking the ratio of twice the diameter to the velocity as an s -state formation time,¹² one has $t_{form} \sim \frac{9n^3}{2\alpha_s^2 M_Q}$ $n = 1, 2, \dots$ and in the ground state $t_{form} \sim 9/2\alpha_s^2 M_Q$. The inverse Bohr radius is the appropriate scale at which to evaluate α_s for the problem, and using the one-loop parameterization for α_s

$$\alpha_s(\mu^2) = \frac{\pi d}{\ell n(\mu^2/\Lambda_{QCD}^2)}, \quad (4.2)$$

where $d = 12/(33 - 2n_f)$, along with $a_0^{-1} = \left(\frac{2}{3}\alpha_s(\mu^2)M_Q\right)$, leads to a condition on $\alpha_s(\mu^2)$:

$$\frac{2}{\pi d} \ell n \left(\frac{2\alpha_s(\mu^2)M_Q}{3\Lambda_{QCD}} \right) = \alpha_s^{-1}(\mu^2) . \quad (4.3)$$

We show a plot of $\alpha_s(1/a_0)$ as a function of M_Q/Λ_{QCD} in Fig. 5. As expected, since the binding scale is much less than the value of M_Q , $\alpha\left(\frac{1}{a_0}\right)$ is significantly larger than $\alpha(M_Q)$. The latter is also displayed for comparison purposes in Fig. 5.

Equating the time of formation and the decay lifetime, one obtains an upper bound on the value of $|V_{Qq}|$ for given values of Λ_{QCD} and M_Q . Choosing $\Lambda_{QCD} = 0.20 GeV$, and $\Lambda_{QCD} = 0.10$ we show $|V_{Qq}|_{max}$ vs. M_Q in Fig. 6.

Fig. 6 clearly shows the point, a surprise to us, that the constraints on V_{Qq} are rather weak, permitting for example $V_{Qq} \simeq 0.3 \sim \sin \theta_c \simeq V_{us}$ for $M_Q \simeq 500 GeV$ with $\Lambda_{QCD} = 0.20 GeV$. As mentioned above, including the Higgs boson, Yukawa coupling effects will only improve chances of bound state formation and further weaken the constraints on $(V_{Qq})_{max}$. This observation is encouraging for prospects of detecting narrow, deeply bound states at higher energies.

Mass splitting between U and D

Turning to the question of mass splitting between the U and D quarks, we use the definition¹³

$$\sin^2 \theta_W \equiv 1 - \frac{M_W^2}{M_Z^2}$$

and the parameterization of radiative corrections in terms of the factor Δr to write¹⁴

$$\frac{M_W^2}{M_Z^2} \left(1 - \frac{M_W^2}{M_Z^2} \right) = \frac{\pi \alpha}{\sqrt{2} G_F} \frac{1}{M_Z^2} \frac{1}{1 - \Delta r} \quad (4.4)$$

The quantity Δr can be expressed as

$$\Delta r = 0.071 - \frac{\cos^2 \theta_W}{\sin^2 \theta_W} \delta \rho_H , \quad (4.5)$$

where the heavy fermion factor $\delta \rho_H$ is given by

$$\delta \rho_H \simeq \frac{3 G_F}{8 \pi^2 \sqrt{2}} \left(M_t^2 + \sum_i \frac{C_i}{3} \Delta M_i^2 \right) , \quad (4.6)$$

and

$$\Delta M^2 = M_1^2 + M_2^2 - \frac{4 M_1^2 M_2^2}{M_1^2 - M_2^2} \ell n \frac{M_1}{M_2} . \quad (4.7)$$

The factor $C_i = 3$ for super heavy quarks and $C_i = 1$ for super-heavy leptons.¹⁵ The masses are designated as M_t for the top quark and M_1 and M_2 for super-heavy fermion generation members. Using the 1992 data book¹⁵ values for M_W and M_Z and the value $\sin^2 \theta_W = 0.230$, we find from Eqs. 4.4–4.7 that the one standard deviation constraint is

$$\Delta M_F^2 \equiv M_t^2 + \sum_i \frac{C_i}{3} \Delta M_i^2 \leq (206 GeV)^2 . \quad (4.8)$$

For the case of one super heavy-quark generation and a degenerate super-heavy lepton generation, the plot of the bound on $|M_U - M_D|$ vs. M_t is shown in Fig. 7. There is evidently room for considerable mass-splitting in the bound (4.8), where $|M_U - M_D| < 140 GeV$ for $M_t = 150 GeV$, for example. The dependence of the bound on the average value of the heavy quark mass is very weak and can be ignored for our purposes.

An independent bound can be obtained from Z decay data. The leptonic and hadronic decay widths, for example, provide useful constraints. Defining the quantity $\sin^2 \bar{\theta}_W \equiv \kappa(M_Z) \sin^2 \theta_W$, one has approximately¹⁶

$$\sin^2 \bar{\theta}_W = \left[\frac{1}{2} - \frac{1}{2} \left(1 - \frac{4\pi\alpha(M_Z)}{\sqrt{2}G_F M_Z^2} \right)^{1/2} \right] \left[1 - \frac{3\alpha}{16\pi \sin^2 \theta_W (1 - 2\sin^2 \theta_W)} \frac{\Delta M_F^2}{M_Z^2} \right] , \quad (4.9)$$

with $\kappa(M_Z) = 1 + \left(3\alpha/16\pi \sin^4 \theta_W \right) \left(\Delta M_F^2/M_Z^2 \right)$. The value of $\alpha(M_Z)$ is obtained from electroweak corrections to α that do not include the heavy quark effects. Using the value $\Delta r = 0.071$ to define $\alpha(M_Z)$ and adopting the first factor in brackets in Eq. (4.9) as the value of $\sin^2 \theta_W$ to be used in the expression in the second bracket, we find

$$\sin^2 \bar{\theta}_W = 0.235 \left(1 - 3.50 \times 10^{-3} \frac{\Delta M_F^2}{M_Z^2} \right) . \quad (4.10)$$

The expressions for the decay widths are

$$\Gamma(Z \rightarrow \ell^+ \ell^-) = \frac{\sqrt{2}G_F M_Z^3}{48\pi} \rho \left[1 + (1 - 4\sin^2 \bar{\theta}_W)^2 \right] \quad (4.11a)$$

and

$$\Gamma(Z \rightarrow \text{hadrons}) = \frac{\sqrt{2}G_F M_Z^3}{8\pi} \rho \left[5 - \frac{28}{3} \sin^2 \bar{\theta}_W + \frac{88}{9} \sin^4 \bar{\theta}_W - \delta_b \right] \times \left(1 + \frac{\alpha_s(M_Z)}{\pi} \right) . \quad (4.11b)$$

The factors ρ and δ_b in Eqs. 4.11 are given by

$$\rho = 1 + \frac{3\alpha}{16\pi \sin^2 \theta_W \cos^2 \theta_W} \frac{\Delta M_F^2}{M_Z^2} ,$$

and $\delta_b = \left[\alpha(1 + 4 \sin^2 \theta_W / 3 - 16 \sin^4 \theta_W / 9) / 8\pi \sin^2 \theta_W \cos^2 \theta_W \right] (M_t^2 / M_Z^2)$, where $\sin^2 \theta_W = 0.235$ is used from Eq. (4.9) onward to evaluate all of the above expressions. The constraints on $|M_U - M_D|$ provided by the 1992 Particle Data Group values of $\Gamma(Z \rightarrow \ell^+ \ell^-)$ and $\Gamma(Z \rightarrow \text{hadrons})$ are illustrated in Fig. 7. We see again that significant mass splitting in a hypothetical fourth generation of fermions is allowed. The constraints on splittings are essentially independent of average mass.

To conclude this section, we note that our assumption of degeneracy between super-heavy generation fermions is consistent with the present bounds, but that there could be shifts in our iso-triplet vs. iso-singlet $Q\bar{Q}$ bound state masses due to ultra-heavy quark non-degeneracy effects. The pattern of a mass-splitting between iso-singlet and iso-triplet states of a given spin and parity would remain, however.

5. Discussion and Conclusion

In the highly relativistic, strong binding regime, we have shown that the contributions of the pseudoscalar, Goldstone-boson degrees of freedom to the binding of ultra-heavy $Q\bar{Q}$ systems play a crucial role in determining the bound state spectrum. Even though these pseudoscalar interactions become negligible compared to the gluon and Higgs exchanges in the weak binding limit, they are solely responsible for the large iso-singlet vs iso-triplet $Q\bar{Q}$ bound state splitting which we found for ultra-heavy, fourth generation, $Q = (U, D)$ quarks. Depending upon the specific approximation scheme used, we saw (Figs. 2,3,4) that the splitting between $I = 0$ and $I = 1$ $Q\bar{Q}$ states become large for M_Q in the range $0.4 \text{ TeV} < M_Q < 0.8 \text{ TeV}$.

Whether a splitting between bound states is large depends of course on the decay width of the (quasi) bound states, and our estimates of $|V_{Qq}|$ in Sec. 4 and the analyses of decay widths of similar states into $q\bar{q}$, H , Z etc. in Ref. 17–21 indicate that the splittings between states will be large compared to the decay widths for the M_Q values in the range given above. The states should, therefore, be clearly separated.

For the production and decay of neutral $Q\bar{Q}$ (and lepton-antilepton) ultra heavy bound states, Refs. 2 and 17–21 provide an encouraging picture of the production and decay signals for such states. The binding indicated by our relativistic, bound state calculations is much stronger than that considered in Refs. 17–18 (gluon non-relativistic potential model only) or 3, 19–21 (Higgs non-relativistic model or Higgs plus gluon non-relativistic potential model).

so the prospects for detecting new heavy bound states, should a fourth generation exist, are made even brighter by our results.

Avenues for further work include the detailed phenomenology of the iso-multiplet system, in particular the possible production and decay of the charged state, and the application of these techniques to TeV scale supersymmetry bound state physics. We plan to address these issues in the future.

Acknowledgements

P.J. and D.M. thank H. Munczek and J. Ralston for useful discussions. This work was performed in part at Ames Laboratory under Contract No. W-7405-Eng-82 with the U.S. Department of Energy and was supported in part by the U.S. Department of Energy Grants Nos. DE-FG02-85ER40214 and DE-FG02-87ER40371, Division of High Energy and Nuclear Physics. One of the authors (A.J.S.) acknowledges financial support from a grant to Iowa State University from the U.S. Department of Education, Graduate Assistance in Areas of National Need Program.

APPENDIX A

We describe here the subtraction procedure used in regulating the B-S equation. Referring to Eqs. (2.1) and (2.2) in the text, we see that the direct, P^2 -pole terms in each equation are independent of q . All of the other terms in the integrands depend on $k - q$ and they decrease like $(k - q)^{-2} \sim q^{-2}$ as $|q| \rightarrow \infty$ for fixed k . Therefore, if the integrals exists, one finds that for fixed P , taking (2.1) in particular,

$$\lim_{|q| \rightarrow \infty} S^{-1}(q_+) \chi_{UU}(q, P) S^{-1}(q_-) \rightarrow \frac{g_2^2}{4} \frac{m^2}{M_W^2} \frac{i}{P^2 - M_Z^2} \gamma_5 \text{Tr} \left(\int \frac{d^4 k}{(2\pi)^4} \chi_{UU}(k, P) \gamma_5 \right) + \frac{g_2^2}{4} \frac{m^2}{M_W^2} \frac{i}{P^2 - M_H^2} \text{Tr} \left(\int \frac{d^4 k}{(2\pi)^4} \chi_{UU}(k, P) \right) . \quad (A.1)$$

If $S^{-1}(q_+)|_{|q| \rightarrow \infty} \rightarrow \not{q}$ as is the case for the perturbative propagators which we use in this paper, then the fixed P , large q limit gives

$$\chi(q, P) \rightarrow \frac{1}{q^2} .$$

In this case, the integral $\int d^4 k \chi(k, P)$ does not exist and the integral equation has no solutions, in accordance with direct numerical studies which we have carried out.

If the q -independent direct (p)-channel pole terms are absent, however, then the q^{-2} dependence of the integrands of all of the other terms in the large q^2 limit means that $\chi(q, P)_{|q| \rightarrow \infty} \rightarrow q^{-4}$ behavior is consistent with the existence of solutions, and we find that this is indeed the case.

To render finite the equations (2.1), (2.2) and their counterparts with U and D interchanged, it is sufficient to subtract each equation at a fixed q , say $q = 0$, because this subtraction removes the q -independent, divergent term. To make this subtraction systematic, one regulates (*i.e.* cuts-off) the integrals, makes the subtraction, and then removes the regularization. The subtracted equation, is as follows:

$$\begin{aligned}
S^{-1}(q_+) \chi_{UU}(q, P) S^{-1}(q_-) - S^{-1}\left(\frac{P}{2}\right) \chi_{UU}(0, P) S^{-1}\left(\frac{-P}{2}\right) \\
= \int \frac{d^4 k}{(2\pi)^4} \left[-\gamma_\mu \chi_{UU}(k, P) \gamma_\nu G^{\mu\nu}(k - q) + \frac{g_2^2}{4} \frac{m^2}{M_W^2} \left(\chi_{UU}(k, P) \frac{i}{(k - q)^2 - M_W^2} \right. \right. \\
\left. \left. + \gamma_5 \chi_{UU}(k, P) \gamma_5 \frac{i}{(k - q)^2 - M_Z^2} + 2\gamma_5 \chi_{DD}(k, P) \gamma_5 \frac{i}{(k - q)^2 - M_W^2} \right) \right] \\
- \int \frac{d^4 k}{(2\pi)^4} \left[-\gamma_\mu \chi_{UU}(k, P) \gamma_\nu G^{\mu\nu}(k) - \frac{g_2^2 m^2}{4M_W^2} \left(\chi_{UU}(k, P) \frac{i}{k^2 - M_H^2} \right. \right. \\
\left. \left. + \gamma_5 \chi_{UU}(k, P) \gamma_5 \frac{i}{k^2 - M_Z^2} + 2\gamma_5 \chi_{DD}(k, P) \gamma_5 \frac{i}{k^2 - M_W^2} \right) \right] .
\end{aligned} \tag{A.2}$$

If the first term on the left hand side of (A.2) is equal to the first integral on the right hand side for each q (*i.e.*, a solution for each q is found), then the companion terms, where $q = 0$, on each side are guaranteed to be equal and the result is an eigenvalue equation for P and q . This equation is, of course, just the original equation without the offending, divergent term and we use this form as our regulated, finite Bethe–Salpeter equation which is to be projected into states of definite quantum numbers.

APPENDIX B

Running of the Yukawa Coupling – The Landau Pole

In this appendix we briefly discuss the effects that arise because of the momentum dependence, or running, of the Yukawa couplings. We work at the one-loop level to see where the Landau pole occurs when the quark masses are in the deep binding region that we probe, $0.5 TeV \lesssim M_Q \lesssim 1.0 TeV$.

The one-loop renormalization group equations for the gluon and Yukawa couplings, g_s and g_Y , without the weak gauge interaction effects, read²²

$$\frac{d}{dt}g_s = -\frac{1}{4\pi^2 d}g_s^3 \quad , \quad (B.1a)$$

and

$$\frac{d}{dt}g_Y = \frac{1}{4\pi^2} \left(\frac{9}{8}g_Y^3 - 2g_s^2 g_Y \right) \quad , \quad (B.1b)$$

where $d = 12/(33 - 4n_g)$, n_g is the number of generations, and $t = \frac{1}{2}\ell n\left(\frac{q^2}{\mu^2}\right)$ with μ an arbitrary renormalization scale.

Parameterizing the solution to (B.1a) in the standard fashion,²³

$$g_s^2(q^2) = \frac{4\pi^2 d}{\ell n\left(q^2/\Lambda_{QCD}^2\right)} \equiv \frac{2\pi^2 d}{t}$$

where $\Lambda_{QCD} \simeq 0.2$ GeV and $\mu = \Lambda_{QCD}$ is adopted, we can rewrite (A.1b) as

$$\frac{dg_Y}{dt} = \frac{9}{32\pi^2}g_Y^3 - \frac{d}{t}g_Y \quad . \quad (B.2)$$

The solution to (B.2) can be written in terms of an integration constant C , to be fixed by the boundary condition:

$$g_Y^2(t) = \frac{1}{Ct^{2d} - \frac{9}{16\pi^2} \frac{t}{1-2d}} \quad . \quad (B.3)$$

Checking several limiting cases of (B.3), we see that

$$g_Y^2(t) = \frac{g_Y^2(0)}{1 - \frac{9}{16\pi^2} g_Y^2(0)t}$$

when $\alpha_{QCD} = 0$ $d = 0$, as it should. The characteristic Landau pole appears at $t = \frac{16\pi^2}{9} \frac{1}{g_Y^2(0)}$. In the weak Yukawa coupling limit, which is the light quark case, where only the g_s^2 effect is kept,

$$g_Y^2(t) = \frac{C^{-1}}{t^{2d}} \quad ,$$

so the quark mass, proportional to g_Y and the weak scale v , behaves as

$$M_Q \sim g_Y(t) \frac{v}{\sqrt{2}} \rightarrow \frac{1}{t^d}, \quad t \rightarrow \infty \quad ,$$

which is the well known leading-log. asymptotic behavior of the light quark masses in QCD.²⁴

The constant C in Eq. (B.3) can be fixed in terms of the input quark mass, M_Q , by requiring the condition²⁵

$$M_Q^2 = g_Y^2 (2M_Q) \frac{v^2}{2} , \quad (B.4)$$

for example. The requirement (B.4) can be solved for C to give

$$C = t_Q^{-2d} \left(\frac{v^2}{2M_Q^2} - \frac{9}{16\pi^2} \frac{1}{2d-1} t_Q \right) , \quad (B.5)$$

with

$$t_Q \equiv \ell n \frac{2M_Q}{\Lambda_{QCD}} .$$

The expression B.3 with C evaluated by (B.5) can now be used to assess the large $|q| > M_Q$, behavior of $g_Y(t)$. We can use the results (B.5) and (B.3) to determine C_Q and q_{pole} , the location of the singularity for various M_Q values. There is no longer a pole at large t , but the coupling still blows up at large t . Interesting values are: $M_Q = 0.5TeV$, where $C_Q = -0.282$ and $q_{pole} = 13TeV$; and $M_Q = 1.0TeV$, where $C_Q = -0.288$ and $q_{pole} = 3.5TeV$. The $t = 0$ pole in (B.3) is an artifact of the standard parameterization of q_s , which is only applicable when $|q| \gg \Lambda_{QCD}$. g_Y should be sensibly constant for small momentum transfers.

The momentum values that are important in the wave function in the Bethe–Salpeter integretion are of the order of the inverse Bohr radius, as a rough rule of thumb. Only in those cases when $M_Q \gtrsim 1TeV$ and in the limit as the bound state mass plunges to zero does the g_Y singularity come near to the relevant range of integration in the Bethe–Salpeter equation. Consequently, the binding energies as a function of M_Q should not be substantially affected by the g_Y singularity below the point where the bound state masses fall to zero, even for the $M_Q \gtrsim 1TeV$ cases.

Figure Captions

Fig. 1: Bethe-Salpeter equations for the $U\bar{D}$ and $U\bar{U}$ bound states.

Fig. 2: Isosinglet and isotriplet masses for mesons in the 0^- channel. Curves S1 (isosinglet) and T1 (isotriplet) are the results from covariant gauge formalism. S2 (isosinglet) and T2 (isotriplet) are the results from Salpeter's equation using only positive frequency components. S3 (isosinglet) and T3 (isotriplet) are the results from Salpeter's equation using both positive and negative frequency components.

Fig. 3: Isosinglet and isotriplet masses for mesons in the 1^- channel. Solid curve (isosinglet) and dotted curve (isotriplet) are the results from covariant gauge formalism. Short dash curve (isosinglet) and long dash curve (isotriplet) are the results from Salpeter's equation using only positive frequency components. Short dash-dot curve (isosinglet) and long dash-dot (isotriplet) are the results from Salpeter's equation using both positive and negative frequency components.

Fig. 4: Isosinglet and isotriplet masses for mesons in the 0^+ channel. Solid curve (isosinglet) and dotted curve (isotriplet) are the results from covariant gauge formalism. Short dash curve (isosinglet) and long dash curve (isotriplet) are the results from Salpeter's equation using only positive frequency components. Dash-dot curve is the isosinglet result from Salpeter's equation with positive and negative frequency components. The isotriplet result for this case is almost identical to the corresponding result obtained by including only the positive frequency components in the Salpeter's equation.

Fig. 5: The value of α_s (solid curve) relevant for the bound state Bethe-Salpeter equation, ignoring higgs interaction, as a function of the quark mass. The dashed curve shows $\alpha_s(M_Q^2)$ which is significantly smaller than the value of α_s that should be used for bound state calculations.

Fig. 6: The maximum value of the CKM matrix element $|V_{Qq}|$ allowed in order for the fourth generation quarks Q to form a meson bound state, q being a lower generation quark. This constraint is derived by including only QCD effects. Inclusion of the Yukawa coupling will allow even higher values of $|V_{Qq}|$.

Fig. 7: The maximum allowed value for the fourth generation quark mass difference $|M_U - M_D|$ as a function of the top quark mass M_t . The three curves represent the constraints using ρ parameter (solid curve), $\Gamma(Z \rightarrow \text{hadrons})$ (short dashed curve) and $\Gamma(Z \rightarrow l^+l^-)$ (long dashed curve).

References and Footnotes

1. Several recent examples are: C. Hill, M. Luty and E. Paschos, *Phys. Rev.* **D43**, 3011 (1991); S. King, *Phys. Lett. B.* **281**, 295 (1992). A recent discussion of ultra heavy quarks and heavy Higgs dynamical effects in a four generation setting is given in T. Truong, *Phys. Rev. Lett.* **70**, 888 (1993).
2. H. Inazawa and T. Morii, *Phys. Lett.* **203B**, 279 (1988). This paper introduces a Yukawa potential, Schrodinger equation version of strong Higgs–boson coupling to fermions.
3. P. Jain, A. Sommerer, D. McKay, J. Spence, J. Vary and B.–L. Young, *Phys. Rev. D* **46**, 4029 (1992). The relativistic bound state formalism used in the present paper is explained in this reference.
4. M. Chanowitz, M. Furman and I. Hinchliffe, *Phys. Lett.* **78B**, 285 (1978); M. Chanowitz, M. Furman and I. Hinchliffe, *Nucl. Phys.* **B153**, 402 (1979).
5. M. Veltman, *Nucl. Phys.* **B123**, 89 (1977); P.Q. Hung, *Phys. Rev. Lett.* **42**, 873 (1979); H. Politzer and S. Wolfram, *Phys. Lett.* **82B**, 242 (1979); **83B**, 421 (E) (1979).
6. We use the term ultra–heavy to distinguish fourth and higher generation fermions from the c , b , and t quarks, generally referred to as heavy quarks.
7. We refer to the $SU(3) \times SU_L(2) \times U(1)$ model with three generations of fermions and one scalar doublet as the standard model.
8. In the unitary gauge, where the Goldstone bosons do not appear, their role is played by the longitudinal degrees of freedom of the $W^\pm$ and Z at the perturbative level.
9. These considerations were not included in Ref. 3.
10. Zero mass bound in strong binding were discussed by J. Goldstone, *Phys. Rev.* **91**, 1516 (1953). For a recent review that includes discussion of this effect, see N. Seto, *Suppl. Prog. Theor. Phys.* **95**, 25 (1988).
11. Vacuum phase transition and tachyonic bound states are studied in scalar theories by B. Haymaker, *Phys. Rev.* **D13**, 968 (1976); *ibid* **D16**, 1211 (1977).

12. I.I. Bigi *et al.*, *Phys. Lett. B* **181**, 157 (1986); J.H. Kühn and P.M. Zerwas, *Phys. Rep.* **167**, 321 (1988); V.S. Fadin and V.A. Khoze, *Pis'mas Zh. Eksp. Theor. Fiz.* **46**, 417 (1987) [*JETP Lett.* **46**, 525 (1987)]; M. Strassler and M. Peskin, *Phys. Rev. D* **43**, 1500 (1991).
13. A. Sirlin, *Phys. Rev. D* **22**, 971 (1980); W. Marciano and A. Sirlin, *Phys. Rev. D* **22**, 2695 (1980).
14. See, for example the discussion of G. Altarelli, Cargèse Summer Institute on Particle Physics 18 July–4 August, 1989 (CERN–TH.5590/89).
15. Review of Particle Properties, *Physical Review D* **45**, Number 11, Part II (1992).
16. See, for example, R. Peccei, *Mod. Phys. Lett. A* **5**, 1001 (1990).
17. V. Barger *et al.* *Phys. Rev. Lett.* **57**, 1672 (1986).
18. V. Barger *et al.* *Phys. Rev.* **D35**, 3366 (1987).
19. H. Inazawa and T. Morii, *Z. Phys. C* **42**, 563 (1989).
20. H. Inazawa, T. Morii and S. Tanaka, *Z. Phys. C* **43**, 569 (1989).
21. H. Inazawa, T. Morii and J. Morishita, *Z. Phys. C* **46**, 273 (1990).
22. C. Hill, *Phys. Rev. D* **24**, 691 (1981). The general framework for Higgs Yukawa couplings was laid out in in T.P. Cheng, E. Eichten and L.F. Li, *Phys. Rev.D* **9**, 2259 (1974).
23. This choice has the advantage of the familiarity. The results that follow are meant to apply at $t > M_Q \gg \Lambda_{QCD}$, however. For an alternative formulation of the solution see, Hill, Ref. 17.
24. H.D. Politzer, *Nucl. Phys.* **B117**, 397 (1976).
25. H. Georgi and H.D. Politzer, *Phys. Rev. D* **14**, 1829 (1976).

Table 1

	V	PS	S
$C, I = 0$	$+\frac{4}{3}\alpha_s$	$\frac{-3}{4\pi} \frac{m^2}{v_0^2}$	$\frac{1}{4\pi} \frac{m^2}{v_0^2}$
$C, I = 1$	$+\frac{4}{3}\alpha_s$	$+\frac{1}{4\pi} \frac{m^2}{v_0^2}$	$+\frac{1}{4\pi} \frac{m^2}{v_0^2}$
Z	$\frac{q^2+q'^2}{2qq'}$	$\frac{q^2+q'^2+M_Z^2}{2qq'}$	$\frac{q^2+q'^2+M_H^2}{2qq'}$

Table 1

Coefficients for the 0^- channel Salpeter equation kernels, Eq. (3.4) and below. Positive signs indicate attractive interactions and the negative sign a repulsive interaction.